



Using Drugs for Pain Prevention: Primary Prevention Across Areas

Introduction

Chronic pain may occur after surgery, trauma, cancer treatment or infection. In the majority of patients enduring chronic pain, a neuropathic or nociplastic component is present which increases pain severity and worsens the quality of life [8]. The transition from acute to chronic pain is a complex process involving multiple biopsychosocial mechanisms. Current knowledge favors mechanisms in relation to both peripheral (at the site of tissue trauma) and central (spinal and supraspinal) sensitization. Drugs have been used to prevent the development of chronic pain, specifically drugs which target spinal excitatory processes and/or which display anti-inflammatory properties able to modulate the release of cytokines from peripheral immune cells and central glial cells.

Drugs for Primary Pain Prevention

The largest bulk of research on the prevention of chronic pain has focused on chronic postsurgical pain, as surgery constitutes one of the common causes of pain and is often scheduled which makes preventive strategies easier to apply. Nonsteroidal anti-inflammatory drugs, gabapentinoids, ketamine and memantine have shown some but inconstant efficacy as preventive analgesics for surgical patients (Please see the IASP fact sheet "Prevention of pain after surgery").

Complex Regional Pain Syndrome type 1 (CRPS-I) is a debilitating chronic pain condition which develops after an inciting event, often minor trauma or surgery, and involves chronic regional pain disproportionate to the causal event. CRPS-I should be distinguished from CRPS type 2, previously called causalgia, where the symptoms are due to nerve damage. The knowledge of the mechanisms of CRPS-I is constantly evolving. CRPS exhibits classic neuropathic pain characteristics but is associated with higher physical disability which considerably lengthens the time of recovery [2]. People who have developed CRPS-I may be at a high risk of recurrence following another trauma or surgical event. Vitamin C supplementation (500 to 1000mg a day for at least 50 days intake) is reported to decrease the risk of developing CRPS type I following wrist fracture and/ or extremity surgery [1, 4]. Vitamin C is a well-documented antioxidant with low risk of toxicity.

Cancer treatment with chemotherapy is known to cause neuropathic pain in 25–50% of patients [5] resulting in an impairment of quality of life. Prevention of chemotherapy-induced pain with anticonvulsants or tricyclic antidepressants has so far not proved very successful [6], but there is some evidence for duloxetine or memantine [7]. Randomized controlled trials and observational studies are needed to achieve clinical significance.

Infection may also have long-standing pain consequences. For example, herpes zoster infection, a painful, blistering skin eruption in a dermatomal distribution caused by reactivation of a latent varicella

zoster virus in the dorsal root ganglia may result in chronic neuropathic pain (post-herpetic neuralgia). Although varicella and zoster vaccines have made major inroads into reducing the burden of disease globally, thus reducing the risk of post-herpetic neuralgia [9], there is currently insufficient evidence to determine the beneficial effect of other antiviral treatments [3].

Conclusion

The prevention of chronic pain remains an area of unmet clinical need. Additional well-designed studies are necessary to determine the overall effectiveness, adverse effects as well as duration of treatment and optimal dosage of preventive drugs. An important step has been done by the inclusion of chronic pain whatever its origin in the current International Classification of Diseases (ICD-11) [8]. Hopefully, this will increase focus on chronic pain and promote research in the field, including the development of preventive strategies.

REFERENCES

- [1] Aïm F, Klouche S, Frison A, Bauer T, Hardy, P. Efficacy of vitamin C in preventing complex regional pain syndrome after wrist fracture : A systematic review and meta-analysis. *Orthopaedics & Traumatology, Surgery & Research: OTSR* 2017;103(3):465-470. <https://doi.org/10.1016/j.otsr.2016.12.021>
- [2] Alam OH, Zaidi B, Pierce J, Moser SE, Hilliard PE, Golmirzaie G, Brummett CM. Phenotypic features of patients with complex regional pain syndrome compared with those with neuropathic pain. *Regional Anesthesia and Pain Medicine*. 2019;44:881-885 <https://doi.org/10.1136/rapm-2019-100511>
- [3] Chen N, Li Q, Yang J, Zhou M, Zhou D, He L. Antiviral treatment for preventing postherpetic neuralgia. *Cochrane Database Syst Rev*. 2014;(2):CD006866.
- [4] Chen S, Roffey DM, Dion CA, Arab A, Wai EK. Effect of Perioperative Vitamin C Supplementation on Postoperative Pain and the Incidence of Chronic Regional Pain Syndrome : A Systematic Review and Meta-Analysis. *The Clinical Journal of Pain* 2016;32(2), 179-185. <https://doi.org/10.1097/AJP.0000000000000218>
- [5] Golan-Vered Y and Pud D. Chemotherapy-induced neuropathic pain and its relation to cluster symptoms in breast cancer patients treated with paclitaxel. *Pain Practice: The Official Journal of World Institute of Pain* 2013;13(1):46-52. <https://doi.org/10.1111/j.1533-2500.2012.00554.x>
- [6] Majithia N, Loprinzi, CL, Smith, TJ. New Practical approaches to chemotherapy-Induced neuropathic pain: prevention, assessment, and treatment. *Oncology (Williston Park)* 2016;30(11):1020-1029.
- [7] Morel V, Joly D, Villatte C, Dubray C, Durando X, Daulhac L, Pickering G. Memantine before Mastectomy Prevents Post-Surgery Pain : A Randomized, Blinded Clinical Trial in Surgical Patients. *PLoS One* 2016;11(4), e0152741. <https://doi.org/10.1371/journal.pone.0152741>
- [8] Treede RD, Rief W, Barke A, Aziz Q, Bennett MI, Benoliel R, Cohen M, Evers S, Finnerup N, First MB, Giamberardino MA, Kaasa S, Korwisi B, Kossek E, Lavand'homme P, Nicholas M, Perrot S, Scholz J, Schug S, Smith BH, Svensson P, Vlaeyen JWS, Wang, SJ. Chronic pain as a symptom or a disease : The IASP Classification of Chronic Pain for the International Classification of Diseases (ICD-11). *Pain* 2019;160(1):19-27. <https://doi.org/10.1097/j.pain.0000000000001384>
- [9] Warren-Gash C, Forbes H, Breuer J. Varicella and herpes zoster vaccine development : Lessons learned. *Expert Review of Vaccines* 2017;16(12):1191-1201. <https://doi.org/10.1080/14760584.2017.1394843>

AUTHORS

Professor Gisèle Pickering, MD, PhD, DPharm,
Clinical Pharmacology Department, CPC/CIC
University Hospital CHU
Clermont-Ferrand, France

Professor Patricia Lavandhomme MD, PhD,
Department of Anesthesiology and Postoperative Pain Service
Cliniques Universitaires St Luc
Université Catholique de Louvain
Brussels, Belgium

REVIEWERS

Rae Bell, MD, PhD
Centre for Pain Management and Palliative Care
Haukeland University Hospital
Bergen, Norway

Christian Duale, MD
Clinical Pharmacology Department, CPC/CIC
University Hospital CHU
Clermont-Ferrand, France

Anne Berquin
Cliniques Universitaires St Luc
Brussels, Belgium

©Copyright 2020 International Association for the Study of Pain. All rights reserved. IASP brings together scientists, clinicians, healthcare providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.

